

Conservation Planning Services and Project Reviews

February & March 2020

**Jason Wagman, Land Use Planning Program
Office of Conservation Planning Services**



Conservation Planning Services

Land Use Planning: 18 staff

- All land/water-use changes in state that have the potential to affect resources (property, activities, species)
- Public and private sources
- Avoidance and Minimization

Landowner Assistance: 18 staff

- Private landowner technical assistance
- Habitat management
- Financial assistance



Conservation Planning Services

- Florida: the FWC is a state review agency and referenced in 38 different rules/statutes
- Federal: USACE, USFWS, DOI, Armed Forces, CZMA/FCMP
- Review projects with or without a federal nexus
- Proactive technical assistance
- Provide permitting guidance



Office of Conservation Planning Services - Land Use Planning - Staff Contacts

Northwest Region

- Bryan Phillips - (850) 767-3646
- Kristal Walsh - (850) 851-8065
- Jennifer Paredes - (850) 617-9408

North Central Region

- Josh Cucinella - (352) 620-7330
- Daniel Penniman - (386) 754-6246
- Kelsey Hamm - (863) 581-8914

Northeast Region

- Laura DiGruttolo - (352) 433-5499
- Michelle Sempsrott - (407) 452-1995

Southwest Region


- Sean Greene - (386) 406-0814
- Jason Wagman - (941) 377-3722 x 6540
- James Keltner - (239) 332-6972 x 9209

Transportation Planning

- Brian Barnett - (772) 579-9746
- Kristee Booth - (850) 363-6298
- Terry Gilbert - (850) 728-1103

South Region

- Christine Raininger - (561) 882-5811
- Vicki Garcia - (561) 882-5711



Administrators: Jason Hight, Program Administrator Ted Hoehn, Project Review Administrator
 Office of Conservation Planning Services - 620 S. Meridian Road, MB 5B5, Tallahassee, FL 32399-1600

FWCConservationPlanningServices@myfwc.com




Conservation Planning Services

- OCPS facilitates this agency function for the FWC
- OCPS also manages the FWC Log that supports this process
- Division/Office lead reviewers
- Land Use Planning is lead on the majority of the reviews
- OCPS Leadership Team



FWC's Project Review Process

- All coordinated through the Conservation Planning Services Leadership Team
- Projects are received through a “general correspondence” inbox
- Projects are screened for appropriateness and logged into the database
- GIS analysis run for each project logged
- Lead reviewer assigned for each project



FWC's Project Review Process

- Technical assistance
 - External - Publicly available information and technical resources, other state and federal agencies
 - Internal – species data, SME collaboration, ISMP
- Identify potential for listed species habitat onsite
- Work on survey guidance and site/project plans WITH applicant – avoidance, minimization, and mitigation
- Requires *constant* internal collaboration to provide an agency opinion



Federal Partners

Federal Agencies - Fish and Wildlife Coordination Act

State Clearinghouse - Coastal Zone Management Act/Florida
Coastal Management Program



US Army Corps of Engineers



FERC



Federal Energy Regulatory Commission



State Partners




- Other State regulatory or review agencies requesting or requiring FWC's review
- Relationships range from a need to know basis to long-standing partnerships



Conservation Planning Services

Growth Management

- LUP coordinates with landowners on ~750,000 acres per year
- Working closely with landowners and local governments on fish and wildlife issues
- Working to keep connections between public conservation lands



Conservation Planning Services

- Local or regional land use changes: Sector Plans
- Commercial/residential/industrial development
- Oil/Gas Exploration: terrestrial and marine geoseismic
- Roadway projects: FDOT and local
- Energy: Sites and corridors
- Mining and reclamation



Stakeholder Engagement

Private Citizens and General Public

- Do we know about it?
- What do we want to do?


Non-governmental or Non-profits

- Formal correspondence
- Discussions in public forums
- Discussions in interagency meetings



Reclamation and Restoration

- Work with large corporate landowners to minimize impacts of their operations
- Recommendations and incentives for reclaiming or restoring the native plant communities
- Mining, power, military, agriculture



FWC Roles

- The FWC is not the agency issuing the permits 95% of the time
- With 34 different rules or statutes mentioning the FWC in state processes - we can always provide information on fish and wildlife resources (almost!)
- We mostly *influence* permitting conditions and recommendations, including mitigation.
- Federal regulations like 33 CFR 332.8(b) REQUIRE the FWC to be involved in interagency review teams.
- Review teams are general practice for many other types of state and local project reviews.



- **Jason Hight, Program Administrator**
Land Use Planning Program
850.413.6966
- **Christine Raininger, Biological Scientist IV**
Land Use Planning Program
561.882.5811



Questions?



Process for FWC Review of FDOT Projects



The FWC is an active participant in several routine transportation project review processes.

- Efficient Transportation Decision Making (ETDM)
- Alternative Corridor Evaluation (ACE)
- Project Development & Environment Study (PD&E)
- Public Conservation Lands Involvement
- Permits



FWC is one of more than 30 members of FDOT's Environmental Technical Advisory Team (ETAT) which includes:

- Federal Agencies
- State Agencies
- Local Governments
- Native American Tribal Governments



Efficient Transportation Decision Making

As part of the Environmental Technical Advisory Team, FWC provides comments on each project run through the ETDM process



How do we review a project, and what elements are most important?

- Most important: the GIS data layers in the Environmental Screening Tool
- Specifically, the data layers in the “Natural “ layer category, which includes the sub-topics:
 - Coastal and Marine
 - Floodplains
 - Geology
 - Land Cover
 - Soils
 - Water Quality and Quantity
 - Wetlands and Surface Waters
 - Wildlife and Habitat



Other routine elements of FWC ETDM reviews

- Aerial and ground-level photography – Google Earth, Street View
- Consult with other FWC staff who are Subject Matter Experts in
 - Fish and wildlife species
 - FWC Management Areas
 - Specific geographic areas (river basins, the Everglades, estuaries, counties, etc.)
 - Agency proprietary interests via fee title ownership or conservation easement
 - Protected species Incidental Take Permits
- Consult with other review agencies, such as the U.S. Fish and Wildlife Service
- Field surveys of selected projects



Draft Memo

Sent to Subject Matter Experts for Review

Final Memo

Sent to FWC official ETAT members

Comments are entered in the ETDM EST



Primary FWC ETDM Project Concerns

- Fish and wildlife habitats that will be impacted, both directly and indirectly
- Fish and wildlife resources, especially listed species, that could be impacted
- Potential for increased habitat fragmentation
- Impacts on wildlife corridors
- Potential for wildlife roadkill



Primary FWC ETDM Project Concerns (cont.)

- Need and appropriate location of wildlife crossings to maintain habitat connectivity and reduce roadkill
- Impacts on public conservation lands, either directly (taken for right-of-way) or indirectly (reduced ability for controlled burning, etc.), with special attention given to impacts on FWC management areas and conservation easements
- Impacts on proposed conservation lands such as Florida Forever or CERP projects
- New public conservation lands that could be acquired as mitigation



Alternative Corridor Evaluation (ACE)

- FDOT identifies multiple feasible corridors for a road project
- An ACE report is prepared and circulated to the ETAT members for review
- Goal is to narrow the list of alternatives for further examination in the PD&E Study
- FWC reviews and compares the alternative corridors using the same criteria as the ETDM process



Project Development & Environment Study

- FDOT's procedure for complying with the National Environmental Policy Act (NEPA) of 1969
- FWC reviews the Natural Resources Evaluation, a part of the PD&E Study that includes a detailed analysis of potential impacts to threatened and endangered species
- For state-funded projects where significant environmental impacts would be expected, a State Environmental Impact Report (SEIR) is prepared, and also reviewed by FWC



Primary FWC NRE Concerns

- Completeness and accuracy of the list of potentially occurring protected species
- Project commitments regarding protected species and their habitats, including pre-construction surveys, further coordination with FWC and USFWS, mitigation for wetlands and species impacts, adherence to species conservation protocols, etc.



Public Conservation Lands Involvement

- If public conservation lands are directly impacted, they must be appropriately replaced
- For state lands, they must be replaced at least at a 1.5 to 1 ratio, with approval of the Governor and Cabinet acting as the Land and Water Adjudicatory Commission FWC is usually involved in the selection process for such mitigation lands
- If the impact is to FWC management areas or conservation easements, we are directly involved in the legal process for mitigation



Permits

FWC Issued Permits

- Most major road projects require FWC gopher tortoise incidental take permits
- For new highways going through lands occupied by several listed species, sometimes a multi-species incidental take permit is required, involving significant acquisition of new conservation lands



Wetland Permits


- All road projects must comply with U.S. Army Corps of Engineers (USACE) permit and state Environmental Resource Permit (ERP) requirements for wetlands and surface water impacts
- USACE permits can involve Section 7 Endangered Species Act Consultation with the U.S. Fish and Wildlife Service for federal listed species, frequently resulting in significant mitigation requirements
- FWC participation in wetlands permit review typically results in FDOT commitments to perform pre-construction surveys for a variety of state-listed species, with requirements for impact avoidance, minimization, and mitigation



Questions?



Florida's Imperiled Species Management Plan & Incidental Take Permitting



Kelly O'Connor
Stakeholder Coordinator, Species Coordination
Kelly.OConnor@myfwc.com
(386) 754-6243

ISMP: comprehensive management plan for state-listed species

- 59 species included
 - 38 state Threatened
 - 21 removed from the List
- Tied to state listing process
- Multiple components
 - Species Action Plans
 - Guidelines
- Rules and Policies
- 10-year plan
- 6 objectives



State-Threatened Species

68A-27.003, F.A.C.

- No person shall *take*, possess, or sell any threatened species, nests, eggs or any part thereof
- Unless authorized by rule, permit, or in a FWC approved management plan or species guideline



State-Threatened Species



American Oystercatcher



Short-tailed Snake



Crystal Darter



Species Action Plan Summaries



Florida Burrowing Owl

Athene cunicularia floridana



Photograph by FWC.

The Florida burrowing owl is geographically distinct from burrowing owls occurring in the western U.S. Burrowing owls require well-drained upland habitats, and are typically found at treeless sites with sparse or low-growing vegetation. Because of this, they may be found at airports, golf courses, in some neighborhoods, and at sites cleared for development.

Status

Listed as state Threatened on Florida's Endangered and Threatened Species List.

Conservation Goal

Conservation status of the Florida burrowing owl is improved to a point that the species can be removed from Florida's Endangered and Threatened Species list and will not again need to be listed.

Identified Threats

- Loss of native habitat and the resulting dependence on altered habitat.
- Lack of protected habitat, even for rural burrowing owls.
- Lure of potentially hazardous habitat, such as predevelopment activities (vegetation clearing) resulting in sites that mimic native habitat of the burrowing owl.
- Urban area threats, including vehicle collision, predation or injury by domestic animals, and burrow destruction by mowers or other equipment.

Conservation Approach


- Protect and manage habitat to support current population and to accommodate population growth.
- Minimize impacts of development and land-use conversion through Species Conservation Measures and Permitting Guidelines, outreach, and technical assistance.
- Ensure adequate protection of burrows through education, enforcement, and management.
- Determine if one or more populations exist, and monitor population(s) to assess size and trend.

Current Protections

- No person shall take (including harm or harass), possess, or sell any Threatened species or parts of their nests or eggs, as outlined in Chapter 68A-27, F.A.C., Rules Relating to Endangered or Threatened Species.
- Under the federal [Migratory Bird Treaty Act](#), it is unlawful to pursue, hunt, take, capture, kill, or sell migratory birds, including their feathers, eggs, and nests.

Resources

- [Species Action Plan](#)
- [Biological Status Review Report](#)



ISMP Policies

- Nest Removal for Inactive Single-Use Nests of State-Threatened Bird Species
- Cryptic Species
- Listed Species and Man-Made structures
- Aversive Conditioning of State-Listed Species



Photo: FWC

Species Conservation Measures and Permitting Guidelines

- Provides information and tools for interpreting existing protections
- Can authorize exempt activities
- Address mitigation options and establishes scientific benefit for state-Threatened species
 - Avoidance
 - Minimization
 - Mitigation



Florida Sandhill Crane

Antigone canadensis pratensis



Photograph by FWC.

Species Overview

Status: Listed as state Threatened on Florida's Endangered and Threatened Species List.

Current Protections

- 68A-27.003(a), F.A.C., No person shall take, possess, or sell any of the endangered or threatened species included in this subsection, or parts thereof or their nests or eggs except as allowed by specific federal or state permit or authorization.
- 68A-27.001(4), F.A.C., Take – to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in such conduct. The term "harm" in the definition of take means an act which actually kills or injures fish or wildlife. Such act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding or sheltering. The term "harass" in the definition of take means an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding or sheltering.
- Florida sandhill cranes, active nests, eggs, and young also are protected under the Federal Migratory Bird Treaty Act, state Rule 68A-16.001, F.A.C., and state Rule 68A-4.001, F.A.C.
- Intentional feeding of sandhill cranes is prohibited under Rule 68A-4.001(5) F.A.C.

Biological Background

This section describes the biological background for this species and provides context for the following sections. It focuses on the habitats that support essential behaviors for the Florida sandhill crane, threats faced by the species, and what constitutes significant disruption of essential behavioral patterns. Florida sandhill cranes (*Antigone canadensis pratensis*) occur from southern Georgia, primarily in the Okefenokee Swamp, to the Everglades (Stys 1997). However, most of the population is in peninsular Florida from Alachua County in the north to the northern edge of the Everglades in the south. The migratory greater sandhill crane (*A. c. tabida*) winters in Florida, arriving in October and November and leaving for breeding grounds in northern U.S. and Canada from late January to early March. Although the two sandhill crane subspecies occurring in Florida are difficult to distinguish, those observed in the peninsula from April to September can be assumed to be the resident Florida subspecies. Florida sandhill cranes typically breed from February through April, but the breeding season can extend as early as December and as late as August (Bent 1926, Walkinshaw 1973). The Florida subspecies and *A. c. tabida* are not known to interbreed.

Habitat features that support essential behavioral patterns

Florida sandhill cranes forage in a variety of open habitats, including shallow (0-32 inches deep) herbaceous wetlands, improved pastures, prairies, open pine forests, croplands, golf courses, airports, and sod farms (Stys 1997). Cranes in north Florida spent 86% of their time in 4 habitat types: pasture, freshwater marsh, pasture-marsh transition, and pasture-forest transition (Nesbitt and Williams 1990). Preferred sandhill crane habitat contains short vegetation (e.g., vegetation less than 20 inches high in uplands), and sandhill cranes generally avoid areas with taller vegetation or dense forest canopies (Stys 1997).

Species Conservation Measures and Permitting Guidelines

- Provides information and tools for interpreting existing protections
- Can authorize exempt activities
- Address mitigation options and establishes scientific benefit for state-Threatened species
 - Avoidance
 - Minimization
 - Mitigation



Osprey *Pandion haliaetus*



Photo by Ricardo Zambrano, FWC.

Species Overview

Status: Removed from Florida's Endangered and Threatened Species list in 2018.

Current Protections

- 68A-4.001, F.A.C., General Prohibitions — No wildlife or freshwater fish or their nests, eggs, young, homes or dens shall be taken, transported, stored, served, bought, sold, or possessed in any manner or quantity at any time except as specifically permitted by these rules nor shall anyone take, poison, store, buy, sell, possess or wantonly or willfully waste the same except as specifically permitted by these rules.
- 68A-1.004, F.A.C., Take—The term shall include taking, attempting to take, pursuing, hunting, molesting, capturing, or killing any wildlife or freshwater fish, or their nests or eggs by any means whether or not such actions result in obtaining possession of such wildlife or freshwater fish or their nests or eggs.
- Osprey, active nests, eggs, and young are also protected under the Federal Migratory Bird Treaty Act and state Rule 68A-16.001, F.A.C.

Biological Background

This section describes the biological background for this species and provides context for the following sections. It focuses on the habitats that support osprey, and the threats faced by the species.

Ospreys occurring in Monroe County are not genetically distinct from ospreys elsewhere in Florida, and this information led to the removal of the osprey of Monroe County from Florida's Endangered and Threatened Species list in 2018. *Pandion haliaetus* species was not previously listed.

The osprey is found in temperate and subtropical areas that contain permanent fresh, brackish, or salt water for foraging (Henny 1988). Ospreys typically utilize canals, ponds, lakes, bays and man-made impoundments. Ospreys prefer areas with clear, shallow waters (0.5 to 2 m [1.6 to 6.6 ft] deep) for hunting that is within 10 to 20 km (6.2 to 12.4 mi) of nest sites. The osprey diet is made up almost entirely of live fish. Bierregaard et al. (2016) report that live fish make up 99% of prey items reported in all publications addressing osprey's diet; ospreys rarely take small birds, mammals, and reptiles.

Ospreys prefer to nest near shallow water but will fly long distances if a breeding site is desirable (Bierregaard et al. 2016). Nests (Figure 1, Figure 2) are large and primarily made up of sticks, historically in trees, but commonly found on artificial structures such as channel markers, utility poles, cell phone



Figure 1: Osprey nest on channel marker in Florida Bay. Photograph by Heather Henkel, USGS.

Stakeholder Involvement & ISMP Resources

- Commenting on draft permitting guidelines
- Feedback on existing Guidelines, policies, etc.
 - Imperiled@myfwc.com or Kelly.Oconnor@myfwc.com
- Additional resources (including Guidelines, Species Action Plans, full text of the ISMP) at <https://myfwc.com/wildlifehabitats/wildlife/>



Incidental Take Permitting



Angela T. Williams
Protected Species Permit Coordinator

Angela.Williams@myFWC.com

(850) 921-5990



How does one “take” a Threatened Species?

- 68A-27.001, F. A. C. - defines **take** of state threatened species as **harass**, **harm**, pursue, hunt, shoot, wound, kill, trap, capture, or collect wildlife
 - **Harm** –activity which directly kills or injures wildlife or indirectly kills or injures wildlife by impairing essential behaviors such as breeding, feeding or sheltering.
 - **Harass** - intentional or negligent act or omission likely to cause injury to wildlife by disrupting normal behaviors such as breeding, feeding or sheltering.



What is Incidental Take?

- 68A-27.001, F.A.C. defines **incidental take** as - any taking otherwise prohibited, if such taking is incidental to, and not the purpose of the carrying out of an otherwise lawful activity.



Permitting Incidental Take

- 68A-27.007, F.A.C.- FWC may issue permits authorizing incidental take of Threatened species if:
 - Species specific standards in Rule 68A-27.003 are met or
 - Clearly enhances the survival potential of the species or
 - Scientific or conservation benefit provided and will not negatively impact survival potential of the species



Permitting Incidental Take (continued)

Considerations for issuing a permit:

1. Objectives of FWC management plan
2. Long range impact
3. Impacts to other fish and wildlife species
4. Extent of injury, harm or loss of the species;
5. Applicant ability to avoid, minimize or mitigate incidental take of the species
6. Human safety and
7. Other conservation and management factors



When is a permit recommended?

Activities will cause take (direct or indirect)

- Avoidance is not feasible (cannot maintain buffer, working during the nesting season)
- Applicants typically meet criteria via combination of
 - Avoidance measures
 - Minimization measures
 - Mitigation measures



Example: Burrowing Owls



- State listed as Threatened
- Prefer grassy habitats with few trees
 - Often occupy residential areas, airports, and sports fields
 - Some dense local populations, such as Marco Island and Cape Coral
- Rely on burrows year round for breeding and sheltering
- Burrows average 5-10 feet long



Example: Burrowing Owls

- Collapsing or blocking a burrow results in take
 - Potential to harm eggs, young, adults
 - Harassment of adults



Example: Beach-nesting birds

- State Threatened
- Nest in “scrape” in sand
- Nests, eggs, chicks camouflaged
- Nest Located: beaches, islands





Example: Beach-nesting Birds

- Person walks through colony, crushes eggs
- Dog runs through colony, birds flush off nest, leaving young and eggs exposed to elements and predators



Example of Take Via Significant Habitat Modification

- Least tern habitat at Summerhaven



When is no permit needed?

- When **take** of species can be completely **avoided**
- ISMP addresses 3 types of avoidance
 - Activities not expected to cause take
 - Species specific in guidelines
 - Avoidance measures undertaken to remove need for ITP
 - Buffers, work during nonbreeding season, JCP
 - Exemptions
 - Land management consistent with management plans
 - Agricultural best management practices
 - Airport Safety Rule – 68A-9.012 F.A.C.



When is no permit needed? (continued)

- When the following ISMP policies are followed:
 - [Nest removal for inactive single-use nests of state-Threatened birds](#) applies to birds that do not re-use the same nests.
 - [Aversive conditioning of state-listed species](#) applies to species with Guidelines that describe non-lethal actions to reduce or eliminate nuisance behavior
 - [State-listed species and man-made structures](#) policy applies to species that are using structures such as buildings, bridges, utility poles, signs and pipes for essential behavior, such as breeding.



Additional Resources

- [Protected Wildlife Permits](#) webpage
- [Species Conservation Measures and Permitting Guidelines](#) webpage
- [Florida's Endangered and Threatened Species List](#)



How to Contact the Permitting Office

(850) 921-5990

or

Wildlifepermits@myfwc.com

or

myfwc.com/license/wildlife/protected-wildlife-permits/



Questions?





Florida Black Bears and Roadways

Bear Management and Research Programs

Florida Fish and Wildlife

Conservation Commission

Background

Abundant

Little or no
management

Over-exploited



Gene Smith



Background

Abundant

Little or no
management

Rare


Rebounding

Recovery
management

Over-exploited

Gene Smith

FLORIDAMEMORY.COM



Background

Abundant

Little or no
management

Conflict
management

Rebounding

Rare

Recovery
management

Over-exploited

Gene Smith



FLORIDAMEMORY.COM

Background




Gene Smith

Over-exploited



FLORIDAMEMORY.COM




ORLANDOSENTINEL




Abundance Estimates

- 2002 Estimate
= 2,600 bears
(Funded by FDOT)
- 2016 Estimate
= 4,050 bears
(> 50% increase)



Bear Range

- Occupied Range
- 49% of historic
- Up from 17% (1993)



Vehicle –Bear Collisions in Florida

- 80% of all known mortality sources (6.1% of statewide population)
- 1976 to 2018: 4,177 mortalities



- Since 2012 > 230 vehicle-bear mortalities/year
- 98.6% are within Frequent or Common ranges


FWC and FDOT Working Together

- Commenting on road development and improvement projects
- Technical assistance and stakeholder groups
- Outreach on driving in bear country
- Periodic assessments of vehicle-bear mortality hotspots
 - Bear Crossing Signs
 - Underpasses
 - Modify Existing Bridges




Vehicle-Bear Mortality Cluster Analysis

- Density Map
- 1km & 5km
 - 2003 to 2007
 - 2008 to 2012
 - 2013 to 2018




Ocala Clusters 2013 to 2018




Fencing Bridges

- FDOT Districts 2,3, and 5
- 2012 to 2016 = 1,238 bears
- 34 bridges identified
 - Proximity to hotspot
 - Ownership on both sides
 - Length and height of bridge







District 5 (Lake Co.) Hotspots and Bridges




Legend




Bear Crossing Sign



Major Roads



County Boundaries




Conservation Areas

Bridges

Priority Level



High




Moderate




Low

1 km Buffer


Density per Sq.km




0 - 0.29




0.3 - 0.96



0.97 - 1.84




1.85 - 3.21




3.22 - 7.44

5 km Buffer

Density per Sq.km




0 - 0.02




0.03 - 0.08



0.09 - 0.19




0.2 - 0.36




0.37 - 0.83


Wildlife Underpass



EXISTING



PLANNED





Bear Crossing Signs




What if I see a Bear?

- 90% of time bears run/climb
- DO NOT run or play dead
- Raise arms, yell, honk horn
- Bear spray



Defensive/Warning Behavior

- Like a rattlesnake
- Personal space
- Stay still, then back away
- If makes contact, fight back





Bat Conservation and Bridges and Other Transportation Structures




Terry J. Doonan, Ph.D.
Jeffery A. Gore, Ph.D.

Florida Fish and Wildlife Conservation Commission

Bat Biology

- The only mammals capable of true flight.
- Wings are supported by bones equivalent to those in our arms and hands.
- Many bats have very long life spans.
- Reproduction is slow for their size.





Bat Biology



- Most bats do not have rabies.
- Many bats hibernate in the winter.
 - But that is rare in Florida.
- In Florida, bats may hibernate just on cold nights/days.
 - Lowers body temperature and metabolic activity.
- Bats can see very well in low light.
- Bats do not try to attack people.




Echolocation



- Bats create ultrasonic calls to navigate and locate prey.
- Calls vary in frequency, pitch, and duration.
- Calls vary depending on a bat's activity.
- Acoustic monitoring equipment is used to identify bats by the frequency and duration of their ultrasonic calls.





White-nose syndrome (WNS)

- A disease that causes the death of hibernating bats.
- Named for the white fungus that infects the bats and appears on their muzzles.
- WNS has caused extensive mortality in cave-roosting bat species across eastern North America.
- Survey results show that WNS has killed more than 6 million bats.
- In many caves, 90 to 100 percent of the bats have died.

Bats in Florida

Florida is home to 13 resident bat species

- Twenty species of bats are found in Florida.
- Only thirteen are resident.
 - 7 considered accidental
- The Florida bonneted bat is found only in Florida.
- Two Resident bat species are listed as Endangered by USFWS:
 - Florida bonneted bat
 - Gray myotis bat



Florida bonneted bat



Legal Status of Bats in Florida

Bats are protected
from take

68A-4.001



Young Brazilian free tailed bat

- Flightless
- Still dependent on its mother.

- There are additional protections during maternity season
 - Maternity season is 15 April -15 August
 - Populations are vulnerable because of loss of young and adults
 - Protects from having dead baby bats in buildings




Nuisance Wildlife Rule: 68A-9.010

- States how and when bats can be removed from buildings.
- Exclusions are the **ONLY** legal removal method.
 - Requires a physical barrier that allows bats to exit and not re-enter
 - Only allowed from August 15th through April 15th
 - Outside the maternity season
 - At least 4 consecutive nights when the projected low temperature is above 50 degrees F.






ALL 3 conditions must be met
for an exclusion to be legal.

Bat Exclusions/Evictions



- Need to be done correctly
- Follow National Guidelines and FWC-UF Recommendations
- Use good materials
- Use appropriate procedures



Bridges Provide Important Roost Sites for Several Bat Species in Florida

5 species known to roost in Florida bridges.

- There are concerns about the Florida bonneted bat

- | | |
|-----------------------|---------------------------------|
| • Free-tailed bat | <i>Tadarida brasiliensis</i> |
| • Big brown bat | <i>Eptesicus fuscus</i> |
| • Evening bat | <i>Nycticeius humeralis</i> |
| • Southeastern myotis | <i>Myotis austroriparius</i> |
| • Big-eared bat | <i>Corynorhinus rafinesquii</i> |
| <hr/> | |
| ?? Bonneted bat | <i>Eumops floridanus</i> |



Brazilian Free-tailed Bat



- Named because tail extends beyond membrane.
- Most common bat in Florida.
- Roosts in many different man-made structures.
- Often found in very large numbers.

Evening Bat



- Dark brown, with muzzle that is unfurred.
- Colonial roosters in man-made and natural structures.
 - Colonies usually from a few to 70 or so bats.
 - Natural roosts include dead trees and woodpecker cavities.
- May roost with Brazilian free-tailed bats.





Big Brown Bat

- Similar in appearance to the evening bat, but slightly larger.
- Also roosts frequently in man-made structures.



Southeastern Myotis

- Small bat.
- Fur is wooly and dull.
- Primarily cave dwelling.
- Can be found in buildings, bridges, and bat houses.
- They sometimes roost in culverts.
- Females form large maternity colonies.



- They preferentially eat mosquitoes!!





Rafinesque's Big-eared Bat

- Enormous ears are distinctive.
- Roosts in small groups, in the open on bridges by hanging on exposed concrete surfaces.
- Rarely encountered in Florida, so status is unknown.



Florida Bonneted Bat



- Florida's largest bat – has a wingspan of 20 inches (50 cm).
- One of the rarest bats in North America – found only in Florida.
- We have limited information about this endangered species.
- Known to roost alone or in small groups.


Potential Issues Associated with the Federally Listed Florida Bonneted Bat

- Within the USFWS consultation area the USFWS typically will ask FDOT to conduct surveys to locate roosts if suitable habitat (including bridges and similar structures) occurs within a project site.
 - Initial inspection mainly is expected to be visual.
 - Where identification of bat presence or bat species ID cannot be made visually, USFWS is likely to ask for acoustic surveys.
- For the construction of new bridges or transportation-infrastructure projects, the USFWS general consultation guidelines would apply.
- The USFWS has draft protocols for conducting roost surveys and acoustic surveys.



-  = State conservation lands
-  = Federal conservation lands
-  = County conservation lands
-  = Private conservation lands

 = USFWS Florida bonneted bat consultation area
- Parts of 14 counties



Potential Issues Associated with the Federally Listed Florida Bonneted Bat

- Construction and maintenance of roads, highways, or bridges may require a consultation with USFWS
- Unauthorized maintenance or repair of bridges or overpasses that are being used as roost sites by the Florida bonneted bat could result in a take – a violation of Federal law.
- They “have been found to roost in tall, live trees or snags; within tree cavities, hollows, and under loose bark. [They also] have been found to use manmade structures (e.g., buildings, bat houses).” In Coral Gables, they have been found roosting in the shafts of royal palm leaves.



Potential Issues Associated with the Federally Listed Florida Bonneted Bat

- The Florida bonneted bat's ability to adapt well to manmade structures may place it at risk from inadvertent or purposeful harm by humans.
- Problems caused by a lack of awareness can cause disturbance or mortality to colonies occupying either manmade structures or natural roost sites.
- To date, the Florida bonneted bat has not been documented using bridges or culverts as roosts.



© Merlin D. Tuttle

Staining and Guano are signs that bats have been, or are, present





Lots of Signs = Lots of Bats

Ft. Pierce, I-95




Brooksville, I-75





Large numbers of bats can
be present in small spaces




There are hundreds of
southeastern myotis
roosting in this 2 inch
wide expansion joint

Where in Florida should you expect bats?

2003 Bridge Bat Survey


301 random bridges


186 non-random bridges



2003 Bridge Bat Survey

152 occupied bridges





Bridge Structure Type

- 1 = Prestressed Concrete Multibeam
- 2 = Prestressed Concrete Slab
- 3 = Steel Stringer
- 4 = Concrete T-beam
- 5 = Prestressed Concrete Continuous Stringer
- 6 = Steel Continuous Girder
- 7 = Steel Continuous Stringer
- 8 = Other Types



Bat Species Identified Roosting in the Bridges

Common Name	Bridges Occupied			Total (%)
	Random	Nonrandom		
Southeastern myotis	0	8	8	(5.3)
Big brown bat	2	33	35	(23.2)
Evening bat	2	8	10	(6.6)
Free-tailed bat	13	91	104	(68.9)






- Bats usually prefer small or narrow spaces in concrete bridges as roosting sites
- Some bats can enter spaces as small as $\frac{1}{2}$ inch wide


A photograph showing a group of dark-colored bats roosting on a light-colored, textured concrete surface. The bats are clustered together in a small group, with some individuals visible in the foreground and others further back. The surface has some visible wear and small holes. A yellow text box is overlaid at the bottom of the image.

Big-eared bats like these typically roost in small groups, in open locations, not in holes or crevices.

A Different Situation


Crystal River, U.S. 19






Hole molded
into the
concrete

- Be aware that culverts can be used as roost sites.
- Bats roost in holes molded into the concrete and in gaps at joints between culvert sections




A cluster of bats in a pipe hole.


- Bats will use many different types of holes or crevices for shelter.



In the wrong spot, bats can cause problems and an exclusion may be necessary

When exclusions have to be done, it is important to have them done well





There may be some limited health considerations for workers:

- Consider wearing masks if working near accumulated guano
- Always wear gloves or use tools to avoid handling dead or injured bats

Recommendations

- Identify existing bat roost sites as early as possible
 - Avoid take issues, contract problems, and work delays
 - Contact FWC for input when necessary
- In South Florida it is important to identify bat species that are present – to avoid take of Florida bonneted bats



A Rafinesque's big-eared bat roosting under a bridge



Recommendations

- Exclude bats only where they create problems for required work
 - Protect them where they are not causing problems
 - Provide workers with information about the bats
- Retain roost sites or provide new roosts whenever possible



Conserving Bats

- Protect natural habitats.
 - Caves, tree roosts.
- More education about bats and their benefits.
- Avoid disturbance.
- If exclusion is necessary, use proper techniques
- Provide alternate roost sites whenever possible.



Box-type Cave Gate Structure



Two double bat houses





QUESTIONS?



Protections for Florida's Birds



Adrienne Doyle

Avian Conservation Biologist

Adrienne.Doyle@MyFWC.com

(352) 620-7732



Craig Faulhaber

Avian Conservation Coordinator

Craig.Faulhaber@MyFWC.com

(352) 620-7346



All Native Florida Birds are Protected


- Even **non-listed** birds are protected
- Some birds have special **federal** protection
- Some birds have special **state** protection



All Native Florida Birds are Protected

- All native birds in Florida are protected by...
 - General prohibitions (68A-4.001 F.A.C.)
 - Federal Migratory Bird Treaty Act (MBTA)
 - State rule adopting MBTA (68A-16.001 F.A.C.)
 - Bird trap rule (68A-16.006 F.A.C.)*

* In effect as of Oct. 3, 2019



Craig Faulhaber



Non-listed Species

68A-4.001, F.A.C., General Prohibitions — No wildlife or freshwater fish or their nests, eggs, young, homes or dens shall be **taken**, transported, stored, served, bought, sold, or possessed...

68A-1.004, F.A.C., Take—The term shall include taking, attempting to take, pursuing, hunting, molesting, capturing, or killing any wildlife...or their nests or eggs by any means...



Non-listed Species

- The **Migratory Bird Treaty Act (MBTA) & associated state rule (68A-16.001)**
 - Prohibits the pursuing, hunting, taking, capturing, possessing, selling, purchasing, & transportation of native birds
 - Also applies to nest, eggs, and parts



Craig Faulhaber



Non-listed Species



Craig Faulhaber

- The **MBTA** formerly covered both intentional and incidental take
- Now interpreted as covering only *intentional* take (M-37050)



Active Nests of Non-listed Birds

- Can *active* nests be taken under MBTA?
 - Yes, if incidental
 - No, if purposeful



Active Nests of Non-listed Birds

- Are active nests protected under state rule?
 - Yes, removal of active nest considered prohibited without state permit
 - Typically, only issued for removal of active nest for human health/safety*



*Would also need federal permit if purposeful



68A-16.006- Prohibitions Related to Bird Traps

- As of October 2019, regulates use & placement, or allowing placement, of bird traps without a permit, with exceptions
- To address illegal trapping of songbirds
- Please report bird traps to FWC law enforcement




Birds with Special Federal Protection



Bald Eagles

- Bald and Golden Eagle Protection Act
- State Eagle Rule (68A-16.002)




The Endangered Species Act

- Federal Law, prohibits “take”
- Definition same as for state Threatened
- USFWS handles permitting
- No separate FWC permit needed



Birds with Special State Protection

- Imperiled Species Management Plan
- [Florida's Endangered and Threatened Species List](#)



State Threatened Species

- 68A-27.003(a), F.A.C. No person shall **take**, possess, or sell any of the endangered or threatened species included in this subsection, or parts thereof or their nests or eggs except as allowed by specific federal or state permit or authorization.





What is
“take?”



What is “take?”

- 68A-27.001: “Take – to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in such conduct.”



What does it mean to “harm?”

- 68A-27.001: “The term ‘harm’...means an act which actually kills or injures fish or wildlife. Such act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding or sheltering.”
- An example: A beach-raker runs over an egg.



What does it mean to “harass?”

- 68A-27.001: “The term ‘harass’...means an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding or sheltering.”
- An example: A beach-raker repeatedly drives through an important breeding site to keep birds from nesting.



But what does take mean
for a particular species?



Permitting Guidelines

- A single source for technical assistance and permitting
- Helps explain what constitutes take for each species
- Available for these birds:
 - burrowing owl
 - sandhill crane
 - osprey
 - white-crowned pigeon
 - wading birds
 - *more on the way!*



Florida Burrowing Owl *Athene cunicularia floridana*



Photograph courtesy Jack Rogers.

Species Overview

Status: Listed as state Threatened on Florida's Endangered and Threatened Species List.

Current Protections

- 68A-27.003(a), F.A.C. No person shall take, possess, or sell any of the endangered or threatened species included in this subsection, or parts thereof or their nests or eggs except as allowed by specific federal or state permit or authorization.
- 68A-27.001(4), F.A.C. Take – to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in such conduct. The term "harm" in the definition of take means an act which actually kills or injures fish or wildlife. Such act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding or sheltering. The term "harass" in the definition of take means an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding or sheltering.
- Florida burrowing owls, active nests, eggs, and young also are protected under the Federal Migratory Bird Treaty Act, state Rule 68A-16.001, F.A.C., and state Rule 68A-4.001, F.A.C.

Biological Background

This section describes the biological background for this species and provides context for the following sections. It focuses on the habitats that support essential behaviors for the Florida burrowing owl, threats faced by the species, and what constitutes take for the species. The Florida burrowing owl is a small, long-legged owl (averaging 9 inches in height) that uses burrows for breeding and sheltering. The Florida burrowing owl's distribution is localized and patchy, occurring primarily in peninsular Florida, with isolated pairs and small colonies found as far west as Eglin Air Force Base and as far south as the Dry Tortugas. The typical breeding season for the Florida burrowing owl is February 15 to July 10, though owls can breed earlier or later. For example, Millsap and Bear (1990) observed egg-laying as early as October 2 and as late as May 9. FWC staff (Zambrano, unpublished data) found that only 4.5% of 89 burrows scoped in south Florida between November 30, 2007, and February 8, 2008, contained eggs or flightless young. Incubation lasts about 4 weeks, and young first start to emerge from the burrow about 2 weeks after hatching (Zarn 1974). Juvenile burrowing owls start learning to fly about 4 weeks after hatching and are starting to fly well by 6 weeks of age (Zarn 1974). Juvenile burrowing owls continue to use their parents' burrows for 30-60 days after they start flying (Mealey 1997).

Habitat features that support essential behavioral patterns

Florida burrowing owls use a breeding burrow and often 1 or more satellite burrows for the essential behaviors of breeding and sheltering. Florida burrowing owls usually dig their own burrows, which are typically 5 to 10 feet long and can be excavated by the owls in as little as 2 days (Millsap 1996). Burrowing owls in Florida are known to use burrows year-round, for roosting during the winter and for raising young during the breeding season (Millsap 1996; R. Mrykalo, personal communication). Some owls leave their burrows for part of the year due to flooding from seasonal rains (Mrykalo 2005). Burrowing owls prefer

Example: State Threatened Wading Birds

- Reddish Egret – Coastal species
- Roseate Spoonbill – Mainly coastal, some inland
- Little Blue Heron and Tricolored Heron – statewide distribution



Photographs (clockwise from top left): little blue heron, reddish egret, and tricolored heron by Jack rogers; roseate spoonbill by FWC.



Example: State Threatened Wading Birds

- Survey seasons vary by region (North, Central, South) and species
- Known breeding sites are re-used and support breeding
 - Need a permit for actions that render part of all of an active or recent breeding site unsuitable for breeding



Photographs (clockwise from top left): little blue heron, reddish egret, and tricolored heron by Jack Rogers; roseate spoonbill by FWC.



Example: State Threatened Wading Birds

Definitions of take:

- Actions that result in loss of habitat within active or recent breeding sites
- Actions that could render a breeding site unusable (e.g. altering water levels, erecting permanent structures within 330 feet of breeding site)
- Ground, aerial, or water based activities that result in repeated flushing of active nests




***Suggested buffer distance =
330 feet from active nests***




Example: State Threatened Wading Birds

- Database of recent breeding sites online:
<http://geodata.myfwc.com/pages/upland> (under Wading Bird Guidelines)
- Not complete, recommend surveys during the breeding season in appropriate habitat



Threatened Wading Birds

- Not all breeding colonies are tree islands
- This wading bird colony in NW Florida is in a ROW that is not flooded year round
- To avoid take when the breeding colony is active, maintain 330-foot buffer



Example: State Threatened Wading Birds

Examples of activities not expected to cause take:

- Changes to canals, ditches or water control structures 330 feet from active breeding sites
- If activities are similar in nature, size, duration, and intensity
- Maintenance of existing roadways within 330 feet of breeding sites, provided outside of the breeding season and no degradation to the breeding site



Example: State Threatened Wading Birds

Other Authorizations:

- Emergency water management actions such as flood control
- Take due to loss of *foraging* habitat for Little Blue Herons, Tricolored Herons, and Roseate Spoonbills will be covered by ERP permits (does not apply to Reddish Egret or nesting sites)



Example: State Threatened Wading Birds

Other Authorizations:

- Routine vegetation maintenance activities within existing highway right of ways that avoid heavy equipment operation within 330 ft of active, known and visibly apparent nests.



Example: State Threatened Wading Birds

Mitigation options:

- Habitat protection or restoration
- Habitat enhancement
- Acquisition, mitigation bank credits or easements
- Funding options
- Information needs
- Programmatic options



Threatened Wading Birds

- Guidelines are available online at <https://myfwc.com/wildlifehabitats/wildlife/species-guidelines/>
- More information on surveys and definitions on take



Little Blue Heron, Reddish Egret, Roseate Spoonbill, Tricolored Heron

Egretta caerulea, *Egretta rufescens*,
Platalea ajaja, *Egretta tricolor*



Photographs (clockwise from top left): little blue heron, reddish egret, and tricolored heron by Jack Rogers; roseate spoonbill by FWC.

Species Overview

Status: The little blue heron (*Egretta caerulea*), reddish egret (*Egretta rufescens*), roseate spoonbill (*Platalea ajaja*), and tricolored heron (*Egretta tricolor*) (hereinafter “wading birds”) are listed as state Threatened on Florida’s Endangered and Threatened Species List.

Current Protections

68A-27.003(2)(a), F.A.C. No person shall take, possess, or sell any of the endangered or threatened species included in this subsection, or parts thereof or their nests or eggs except as allowed by specific federal or state permit or authorization.

68A-27.001(4), F.A.C. Take – to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in such conduct. The term “harm” in the definition of take means an act which actually kills or injures fish or wildlife. Such act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding or sheltering. The term “harass” in the definition of take means an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding or sheltering.

State-Threatened wading birds, active nests, eggs, and young also are protected under the Federal Migratory Bird Treaty Act, Rule 68A-16.001, F.A.C., and Rule 68A-4.001, F.A.C.

Biological Background

Florida’s Endangered and Threatened Species List includes 4 state-Threatened wading birds: the little blue heron, reddish egret, roseate spoonbill, and tricolored heron. The snowy egret (*Egretta thula*) and white ibis (*Eudocimus albus*) were removed from Florida’s Endangered and Threatened Species List in 2017 and will be discussed in a separate set of Guidelines. This section describes the biological background for the 4 state-Threatened wading birds (hereafter, wading birds) and provides context for the following sections. This section focuses on the habitats that support essential behaviors for these species, threats faced by the species, and what constitutes take for the species. For more information about life history and conservation actions for wading birds, please refer to [A Species Action Plan for Six Imperiled Wading Birds](#).

All 4 species are year-round residents in Florida (Greenlaw et al. 2014). Little blue herons, roseate spoonbills, and tricolored herons are found throughout the state, though roseate spoonbills currently nest only in peninsular Florida (Greenlaw et al. 2014; Florida Ornithological Society, unpublished data). Reddish egrets are found almost exclusively in coastal areas, with nesting occurring from south Florida to at least Levy County on the Gulf Coast and Brevard County on the Atlantic Coast (Lowther and Paul 2002, Cox et al. 2017b).

QUESTIONS?





Sea Turtles and Lights

Balancing Property Rights, Safety,
and Sea Turtle Survival



Tonya Long, Luke Davis, Rachael Stevenson
Imperiled Species Management
Florida Fish & Wildlife Conservation Commission

Sea Turtles Nesting in Florida



Photo: M. Nicholas, Gulf Islands National Seashore, MTP #032

★ **Loggerhead**
(*Caretta caretta*)
Status: **Threatened**



Photo: FWC

★ **Leatherback**
(*Dermochelys coriacea*)
Status: **Endangered**



Photo: B. Witherington, FWC

★ **Green**
(*Chelonia mydas*)
Status: **Threatened**



Photo: T. Hirama, FWC

Kemp's Ridley
(*Lepidochelys kempii*)
Status: **Endangered**



Photo: N. Brockhoff

Hawksbill
(*Eretmochelys imbricata*)
Status: **Endangered**



Sea Turtle Nesting 101

- Nesting season is May 1 – October 31 (March 1 – October 31 in Brevard through Broward)
- Nesting and hatching both occur at night.
- Nests contain ~100 eggs and incubate ~2 months
- Females return to nest in the same areas where they hatched.
- About 40% of worldwide loggerhead nesting occurs in Florida – more than anywhere else in the world!





Photo: T. Long, FWC


Why is artificial lighting such a problem?

Hatchlings rely on multiple visual cues to find the water

- TOWARD bright open areas/horizons
- AWAY from dark silhouettes i.e. dunes and vegetation
- Attracted to SHORT wavelength light
- Less disturbed by LONG wavelength light




Witherington et al. 2014





Conflicting cues often lead to disorientation.



Undisturbed
hatchlings:
seaward orientation
in a “V” formation



Disoriented
hatchlings:
circular orientation or
not towards the ocean




Nesting females can disorient too!



Tracking Disorientations

www.MyFWC.com

Search “disorientation”



Florida Fish and Wildlife
Conservation Commission

Site Search



Home > Wildlife & Habitats > Wildlife Conservation > Sea Turtle Program > Sea Turtles and Lights > Sea Turtle Disorientations

Sea Turtle Disorientations

Disorientations and Hazards to Sea Turtles

Expand All

Collapse All

What are disorientations?



Disorientation Hazards: Adult Nesting Females



Disorientation Hazards: Hatchlings




Get Involved



Local Lighting Ordinances



Education, Information, & Kids Activities



Disorientation Photo Gallery

Disorientation Data

Review the PDF charts for the past five years of data for each species or follow the Disorientation Map action tab for a geographical representation of disorientation reports throughout Florida's coastline.

[Loggerhead Disorientations.pdf](#)

[Green Turtle Disorientations.pdf](#)

[Leatherback Disorientations.pdf](#)

[DISORIENTATION MAP](#)



NATIONAL FISH AND WILDLIFE FOUNDATION GRANT

Sea Turtle Disorientation Data Collection and
Management

GRANT DETAILS

You've identified the problems – now what?

Lights should follow
FWC's Three Golden Rules:


- 1) Keep it Low
- 2) Keep it Long
- 3) Keep it Shielded



FWC Golden Rule Number 1

Keep it Low

- Keep mounting heights low to minimize light trespass.
- Keep it low also refers to lumens. Use only the lumen output needed. For most applications this can be minimal.




FWC Golden Rule Number 2

Keep it Low

Keep it Long (wavelength)

Use bulbs with wavelengths greater than 560 nm.



FWC Golden Rule Number 3

Keep it Low

Keep it Long

Keep it Shielded

Eliminate the point source of light.

- Use full cut-off or fully recessed fixtures.
- Use 180 to 270 degree external beachside shields when in proximity to the beach.



Can streetlights really meet the Golden Rules?

YES!

Synergy Streetlight demo with FDOT –
now being installed in Fort Lauderdale

Cree RSW Series Streetlight demo with
FPL – now on FPL approved products list



Photo: T. Long, FWC



Photo: T. Long, FWC

FDOT Wildlife-Sensitive Roadway Designs

PROSECUTION AND PROGRESS – LIMITATIONS OF OPERATIONS – NIGHT WORK ALONG COASTAL ROADS. (REV 4-8-19) (FA 7-22-19) (1-2020)

SUBARTICLE 8-4.1 is expanded by the following new Subarticle:

- **8-4.1.1** Additional Requirements for Night Work Along Coastal Roads: The project is located adjacent to sea turtle nesting habitat. Direct all work zone lighting away from the beach to avoid illumination of or direct visibility from the beach. Shield luminaires to avoid lighting areas outside of the immediate construction area.

FDOT Design Manual (1/2020)

231.2.1 Wildlife-Sensitive Conventional Lighting

For conventional lighting near a wildlife area of concern (as determined by the Office of Environmental Management), incorporate the following design requirements:

- 1) Where feasible, orient luminaires away from the wildlife area of concern.
- 2) Design lighting system using luminaires from the Wildlife-Sensitive Conventional Lighting category of the APL.
- 3) Use the criteria for Wildlife-Sensitive Conventional Lighting from Table 231.2.1 in accordance with the requirements of FDM 231.3.

Standard Specifications for Road and Bridge Construction (1/2020)

992-2.4.2 Luminaires for Wildlife-Sensitive Conventional Lighting: Luminaires must meet the following additional requirements:

- a. The light source for the luminaires must be true red, orange, or amber light-emitting diodes (LEDs) with no more than 1.75% of the spectral power distribution below 560 nm. Submit testing report.
- b. The optics must have an IP 66 rating. Submit testing report.
- c. The luminaire mounting assembly must be a slipfitter type designed to accommodate a nominal 2 inch pipe size (2-3/8 inch O.D.) arm or a pole top mounting assembly designed to accommodate a 2-3/8 inch pole top tenon.
- d. Luminaires must have a IESNA light distribution curve (IES LM-79) designated by an EPA-recognized laboratory. Submit testing report.
- e. Luminaires must meet a minimum pole spacing of 50 feet using the AGi32 lighting optimization tool in accordance with the settings shown in 992-2.4. Submit IES file.

FLORIDA DEPARTMENT OF TRANSPORTATION



STANDARD SPECIFICATIONS FOR ROAD AND BRIDGE CONSTRUCTION JANUARY 2020

“Halfway” Solutions to Lighting Problems

Timers, switching off lights, or changing LED colors seasonally are NOT recommended long-term solutions.

But WHY?

- Prone to malfunctions
- Turning off lights may create unsafe situations for drivers/pedestrians
- Switching colors fosters the perception that amber lighting is “less safe” than white lighting



Photos: T. Long, FWC



Thank you!
For questions, contact
SeaTurtleLighting@MyFWC.com

Want more info?
www.MyFWC.com/seaturtle

Wildlife Crossings and Passage Structures

February 2020



Sean Greene, Land Use Planning Program
Office of Conservation Planning Services



Wildlife Passage Structures


- Three primary types:
 - Bridge underpasses
 - Culverts
 - Bridge overpasses
- Most originally designed to pass water beneath road infrastructure
- Fencing must be used to direct wildlife to passage structures
- Tend to focus on large mammalian species



amusingplanet.com



humansandnature.org




arc-solutions.org



Wildlife Crossings in Florida

- Range in size from 12 inch culvert pipes to the bridge underpasses on I-75 that measure 8 feet high and 120 feet wide
- First wildlife crossings were included during the planning for the conversion of Alligator Alley to I-75 in 1972
- Nearly 100 wildlife crossings in Florida today (mostly in South Florida)
- FDOT, FWC, and USFWS consider the need for wildlife crossings in all new road construction and expansion



Wildlife Vehicle Collisions (WVCs)

- Reduce Wildlife Vehicle Collisions (WVCs)
 - 2 million WVCs per year
 - 29,000 human injuries
 - 200 human deaths
 - \$8.4 billion in damages
 - 8-20% increase per year
- Florida Black Bears
 - WVCs account for over 90% of bear deaths in FL
 - WVCs with bears have increased steadily since 1981
 - Over 50% increase in last 10 years




K. Gunson, G. Mountrakis, L. Quackenbush (2011)



Florida black bears can weigh up to 740 pounds.



Florida Panther



- 1981-1999: 1.5 mortalities per year
- 2000-2017: 15+ mortalities per year
- Early measures:
 - Lower speed limits at night
 - Widening road shoulders
 - Reflectors
 - Rumble strips
 - Public information
- Most effective (and most expensive) measure remains wildlife crossings
- Early measures reduce WVCs, wildlife crossings can eliminate them



Wildlife Populations

- Habitat Connectivity

- Deer – 7 square miles
- Black Bear – 15 square miles
- Coyote – 60 square miles
- Panther – 200 square miles

- Genetic Isolation

- “Island Populations”
- Increased inbreeding and disease risk
- Approximately 20 black bears isolated at Chassahowitzka





Illustration by Wildlands Network



Case Study: Lake Jackson Ecopassage

- 4000 acre sinkhole lake bisected by US 27
- Turtles regularly cross in search of water
 - 2,710 crossing attempts per year per mile
- 11,000+ WVCs over 5 year period
 - 98% mortality rate for crossings
- FDOT, Capital Regional Transportation, Leon County, and NGOs work to address issue
- Construction of solid wall and multiple culverts in 2010
 - \$3.4 million in federal stimulus money
 - 0 recorded mortalities since




Lake Jackson Turtle Rescue



Small Animal Design

- Entrance coverage
 - Especially for prey species
 - 2013 Montana Study
 - 42.9% increase in passage rate when coverage added
- Shelving throughout crossing
 - Especially when water present
 - 2013 Montana Study
 - 0 small animal species use of wet culvert without shelving
 - 14 species once shelving added



Use of Elevated Paths



Large Animal Design

- Require a large cross-sectional space (passage height x width)
 - Limited to large (7+ foot diameter) culverts or bridge underpasses
- More sensitive to influence of human co-use
 - Large mammals will develop strategies to minimize human interactions
 - Temporal divergence (humans at day, wildlife at night)
 - Physical divergence (humans cross on north path, wildlife on south)
- In Washington study, one side of bridge underpass had a cement trail frequented by humans during the day
 - Over 90% of wildlife crossings occurred on the opposite side



Conclusions



- Bridge underpasses and culverts play a valuable role in maintaining habitat connectivity and reducing wildlife vehicle collisions
- Wildlife crossings are the only method proven to fully eliminate wildlife vehicle collisions on targeted sections of roadway
- A wide variety of species will make use of wildlife crossings
- Specific design elements can increase the utility of wildlife passage structures for specific types of species



- **Jason Hight, Program Administrator**
Land Use Planning Program Administrator
(850) 413-6966
- **Sean Greene, Land Use Planning Biologist**
14495 Harllee Road
Palmetto, FL 34221
(386) 406-0814



Questions?





Gopher Tortoise

Biology, Conservation, and Regulation

FDOT - 2020



Gopher Tortoise (*Gopherus polyphemus*)

- Only tortoise in North America east of Mississippi River
- Candidate for Federal Listing (ESA)



Gopher Tortoise (*Gopherus polyphemus*)

- Live to 40-60 years old
- Adults average 10 inches long
- Slow to reach sexual maturity
 - Between ages of 9 to 21



Adults



Juveniles



Nests

- High predation on nests/juveniles
- Juvenile's shell stays soft for up to 7 years



Diet

- Herbivores
 - Low-lying vegetation
 - Broadleaf grasses, gopher apple, prickly pear, cactus, berries, and more
- Opportunistic
- Go up to 160 ft for food



Behavior & Home Range

- Active 60° to 95° F
 - Foraging, burrowing
- Bask in sun to increase body temp
- Stay close to burrow, except on foraging journeys or to mate
- Males tend to have a larger range



Adult Females	Adult Males	Juveniles (<4 yrs)
0.2-1.4 ac	1.1-3.2 ac	0.02-0.9 ac



Habitat

Scrub



Longleaf pine

Coastal dunes



Pine flatwoods



Habitat



Burrows

- Average 7 ft deep and 15 ft long
 - Up to 40 ft long
- Can have more than 1 burrow within home range
- Half-moon shape



Burrows

- Proportional to size of tortoise
- Spend 80% of life in burrow
- Refresh burrow in fall and rest in winter





Burrows



Keystone Species

KEYSTONE SPECIES



Identifying GT Burrows

- Potentially Occupied
 - Active
 - Inactive
- Abandoned



Potentially Occupied

- Active
 - Half-moon shaped entrance
 - Clear of vegetation
 - Good repair
 - Loose soil
 - Tortoise tracks



Potentially Occupied

- Inactive
 - Half-moon shaped entrance
 - Vegetation growing
 - Good repair
 - Hard-packed soil
 - No tortoise tracks
 - Partially covered with leaves




Abandoned

- Appears unused or dilapidated
- Entrance partially or completely collapsed
- Filled partially or completely with leaves or soil



Why are populations declining?

- Habitat loss
- Fragmentation
- Degradation
- Disease
- Illegal harvest



Why are populations declining?



Upper Respiratory Tract Disease (URTD)

- Symptoms = discharge from nasal passage or eyes, swelling of area around eyes



Illegal Harvesting



What Actions Are Being Taken?

- Permitting
- Local Gov't Coordination
- Law Enforcement
- Habitat Preservation
- Habitat Management
- Population Management
- Disease Management
- Incentives
- Monitoring
- Education and Outreach
- Future Research



Photo Courtesy of www.floridastateparks.com



Management Plan (approved 2012)

- Goal = Restore/maintain viable populations so GT no longer needs listing
- Conservation action = regulation, permitting, law enforcement, etc.

Permitting Guidelines

- Permits provide authorization to impact tortoises and their burrows



Protections

- F.A.C. 68A-27.003(2)(d)3

No person shall:

- take or attempt to take
- pursue, hunt, harass, or capture
- possess, sell, or transport any gopher tortoise or parts thereof or their eggs
- OR molest, damage, or destroy gopher tortoise burrows



Burrow Protections

- Without a permit, no person shall:
 - Place harmful substances or devices inside a burrow
 - Collapse any part
 - Block, cover, or fill
 - Exclude tortoises



Activities that **Do Not** Require a Permit

Bona fide...

- Routine agricultural practices
- Silvicultural activities
- Wildlife management activities
- Linear utility and highway right-of-way vegetation maintenance
- Routine yard maintenance that **does not damage a burrow**
- Hand trimming vegetation



Tips for Vegetation Maintenance

- Do not stand/drive equipment directly on top
- Mow at least 5 ft radius around burrow
- Hand trim up to burrow entrance
- Flag/stake hard-to-see burrows
- Use pesticides and herbicides according to label instructions + not immediately around burrow entrance



Activities that **Do** Require a Permit

- Any activity that may cause take, harassment, or molestation of GTs, or destruction of their burrows:
 - Clearing, grading, paving
 - Bulldozing, digging
 - Construction
 - Site preparation for development



Permit Types

- Authorized GT Agent
- Relocation
 - 10 or Fewer Burrows
 - Conservation
 - Temporary Exclusion
 - Disturbed Site
- Recipient Site
 - Long-term protected
 - Short-term
 - Public Conservation Lands
- Burrow Structure
- Scientific Collection




Authorized GT Agent Permit

- Required when relocating tortoises to off-site location, for temporary exclusion permits, and for disturbed site permits
- Not all agents are permitted to perform every task



10 or Fewer Burrows Permit


- Projects with no more than 10 burrows impacted



Conservation Permit

■ F

ted




Disturbed Site Permit

- Site prep activities occur before GTs are moved



How to Avoid Need for a Permit

- Maintain 25-ft radius buffer during ground disturbance (e.g., fence installation, clearing vegetation or construction)



Avoid Creating “Islands”/“Crop Circles”



Tortoise-wise Community Program

- Tortoises do live in neighborhoods!
- Landscape with native plant species
 - Florida Guide to Gopher Tortoise Friendly Plants
 - Available at [MyFWC.com/GopherTortoise](https://myfwc.com/gophertortoise)
- Keep dogs away from burrows and tortoises
- Help tortoises across roadways
- Download GT App
- Report mortalities



Tortoises and Roads

- Move it to other side
→ in direction it was headed
- If injured: Contact Assistant Biologist to locate Rehabber (850) 921-1030



Tortoises and Dogs

- Owner is legally responsible for dogs
- Contact Assistant Biologist for options
(850) 921-1030



Waif Tortoises

- Waif = GT from unknown location, or kept as pets for many years
- Contact Assistant Biologist for options
[\(850\) 921-1030](tel:8509211030)
- Please **DO NOT** relocate to parks, etc.
→ possible disease transmission!



Online Access to Permits


- Public online map shows issued permits

MyFWC.com/GopherTortoise/Permitting



Online Permit Locator Map

<https://public.myfwc.com/maps/gtmapping/default.aspx>





Wildlife Alert Hotline

- Wildlife Alert Hotline: 888-404-FWCC (3922)
- Tip Line: Text or email Tip@MyFWC.com
- Wildlife violations vs Permit violations



Tortoise App




<https://myfwc.com/wildlifehabitats/wildlife/gopher-tortoise/app/>

Report Mortalities

Gopher Tortoise Mortality

To better protect and understand the gopher tortoise, data on mortality observations in Florida can help track the density of gopher tortoises in urban areas. The FWC and other conservation groups can use this data to further protection efforts. Florida's citizens are a valuable resource when it comes to gathering data. This survey will help to determine areas of higher mortality and help the FWC and other groups develop new plans to enhance the protection of the gopher tortoise.

[Read More Information](#)



Dead Gopher Tortoise Form

[Read Instructions](#)

* Required fields

Latitude *

27.990844851725232

Longitude *

-81.55570080895947

Date of Sighting *

(MM/DD/YYYY)

Carapace Size (Upper shell length)

☐ 1-7 inches

☐ > 7 inches

☐ Unknown

Best Description of Me

☐ Professional (biologist, forester, professor)

☐ General Public

☐ Other

My Name *

Organization Name


Email *

<http://MyFWC.com/GopherTortoise>



Regional GT Biologists

- SW: Kyle Brown
- S (+ Brevard, Indian River, Orange, Osceola): Momoka Maeda
- NE: Samantha Cobble
- NW/NC: Eric Seckinger



THANK YOU!



<http://MyFWC.com/GopherTortoise>





Wildlife Assistance Program

Rhonda Douthett

FL Fish and Wildlife Conservation Commission

Wildlife Assistance Biologist



Current trends

- Human-wildlife conflict & Floridians
- Increase in overall complaints
- Public not aware of agency role
 - Media covers trapping & rescue events of:
 - Sea turtles
 - Panthers
 - Manatees
 - Bears




Florida Panther, FWC photo- Palm Beach County



Florida black bear, FWC photo- Lake County


Mission: Minimize adverse impacts of fish and wildlife on Florida's environment, economy, and human health and safety.



Wildlife Assistance Program



...Addressing Conflict, Fostering Coexistence

- Six regional staff providing technical assistance to foster coexistence
 - Primarily over the phone
 - Equipment loans available
- Partnerships and coordination with other agencies and NGOs
- Pro-active engagement of communities and neighborhoods




Wildlife Incident Management System (WIMS)

- Statewide database for recording incidents regarding conflict wildlife, wildlife assistance, and nonnative wildlife
- Maps incidents for visual to help pinpoint areas for proactive outreach.
- Imports LE CAD calls for bears, panthers and coyotes daily for review and follow-up if needed




Wildlife Assistance Program Totals



Calls by Native Species Type, Fiscal Year 2017-2018 (non-bear)

Total = 5,845



Coyote biology

- Omnivorous, highly adaptable
- Adults 25-40 lbs.
- Home range size:
natural areas ~15 mi²,
urban areas ~3 mi²
- Mate in Winter
- Pups in Spring
- Litter size ~6 pups
- Both parents care for
young



FWC photo- Clearwater



FWC trail camera photo- Big Cypress



Coyotes in Florida: the Ugly?

Coyotes have an undeserved public reputation

Common myths and misconceptions:


- Size
- Packs
- Coy-dogs/ Coy-wolves
- Disease
- Hollywood
 - Wiley coyote



FWC photo- Clearwater



Actual weights of coyotes




FWC diet study-149 coyote carcasses
collected statewide from 2011-2013



FWC Coyote-Related Incidents Point Density
Fiscal Year 2013-2018

● Outreach Presentation



0 25 50 100 Miles



Raccoon biology



- Eat a wide variety of prey:
 - Fruits, vegetables acorns, seeds
 - Eggs
 - Crayfish, frogs, small mammals
 - Fish
 - Human-related foods
- Typically nocturnal, but day activity not unusual
- *Densities of 100 per square mile can be attained with abundant food resources*
- Breeding occurs at one year of age, with litters born in March and April
- Litters average 3-4 young



Wild Hog (*Sus scrofa*)

- Nonnative
- Diet- primarily vegetation, but will eat just about anything
- Damage- feeding on crops, tusking & rooting, wallowing, disease



FWC Photo



Wild Hog Range

- 500,000+ in Florida (in every county)
- 1-2,000,000 in the Southeast
- Several million in 39 states and provinces
- ~1 mile² (450-740 acres)
 - Depends on food
- 1 hog / 32 acres



Wild Hog Biology

- Start breeding at 1 yr of age
- Breeding peaks in spring & fall
 - Produce 2 litters of 1-13 piglets each
- Full grown in 3-5 years
- Average life span: 4-5 years
- Solutions
 - Exclusionary fencing
 - Shooting/ hunting
 - Trapping



Panther vs. Bobcat



Identifying Florida's Native Cats



Florida Panther



Description: uniformly tan, adults not spotted, tail nearly length of body

Weight: 60-160 lbs

Total length: 7-8 ft


Body Length: 4.5 ft

Tail Length: 3 ft

Shoulder height: 2.25 ft

Back of ears: black

Tip of tail: black all around



Bobcat



Description: reddish brown, spots evident but variable, tail much shorter than length of body

Weight: 20-30 lbs

Total length: 3 ft

Body Length: 2.5 ft

Tail Length: 6 in

Shoulder height: 1.5 ft

Back of ears: white spot

Tip of tail: white underside



Reporting Panther Sightings:



Panther Sightings

Florida Panther

Panthers naturally occur in Florida. Help FWC improve our knowledge of panther distribution in Florida and revise FWC's range map.

[Want to see the sightings or more photos?](#)


[View Panther Sightings Map](#)

[Read More...](#)



Instructions

- Drag the red marker  to the approximate location and then zoom into the map to help pin-point the desired location. Alternatively, you can enter your lat/long in form below. Accepted lat/long format is dd.dddd or -dd.dddd.
- Complete the form with the sighting date, your name, and contact email.
- Optionally attach up to three images using the Plus button below the form. When finished click the Submit button.



Panther Sighting Form

Latitude *	Longitude *
Date of Sighting *	



Burmese python



Dark arrowhead-shaped wedge on back of head

Dark irregular blotches on tan background (like puzzle pieces that fit together or the markings on a giraffe)



Burmese pythons/Photo credit Edward Mercer

Burmese Python



6/4/2014, Cuthbert Lake Wood Stork Nesting Colony



Burmese Python Observations and Removals

- Removals
- Found Dead
- Observation



Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroX, Geomapping, AeroGRID, IGN, IGP, swisstopo, and the GIS User Community



Argentine black and white tegu

Thick jowls on
males

White bands across
dark body



Argentine black and white tegu/Photo credit FWC



Tegu



Black and White Tegus Observations and Removals

- Removals
- Found Dead
- Observations

Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AEX, Getmapping, Aeragrid, IGN, IGP, swisstopo, and the GIS User Community



Boa constrictor



Tan ovals separated by dark brown saddles; ovals turn red near the tail on red-tailed boas


Thin dark-colored line on head



Boa constrictor/Photo credit Adam Stern

Boa Constrictors Observations and Removals

- Removals
- Found Dead
- Observations



Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AEX, Getmapping, Aergrid, IGN, IGP, swisstopo, and the GIS User Community



Ball python



Light dumbbell-shaped
(or alien head-shaped)
blotches on dark
background

Dark wedge on back of
the head, like Burmese
pythons

Ball python/Photo credit Adam Stern



Nile monitor





Bands of spots or
“coins” across
back

Bands across tail



Nile monitor





Seen something exotic? Report your sighting!

- 1. Take a picture**
- 2. Note the location**
- 3. Report your sighting**

By phone: 888-Ive-Got1 (888-483-4681) **Online:** IveGot1.org
or download the IveGot1 reporting app.

Knowing the distribution of nonnative species in Florida helps wildlife biologists plan more effective management strategies.

MyFWC.com/Nonnatives



Native snakes



Red Rat Snake



Water Snake



Water Moccasin



Safety First

- Always use caution
- Park a short distance away
- Take photos while approaching
- Keep a distance of 15 ft.




Who should you contact when there is a problem?

- FWC SW Regional Office (863) 648-3200
- SNAP: (866) FWC GATOR
- Nuisance Wildlife Trapper
<http://myfwc.com/trappers/>



FLORIDA FISH AND WILDLIFE CONSERVATION COMMISSION
HEADQUARTERS AND REGIONAL OFFICES



Suggestions

- Violations can be reported to the Wildlife Alert Hotline
- **1-888-404-3922**
- Reports can be made anonymously

