On SR85, Eglin Parkway, adjacent to Eglin Air Force Base in Okaloosa County is a naturally occurring wildflower area that, by request of the County Commission, shall be managed for native wildflowers along the southbound side of the right of way for approximately 1.25 miles. Bordered by a Coastal Scrub Forest of mixed pines and oaks, dominated by Sand Pine and Laurel Oak, the native understory is a mix of shrubs and forbs adapted to fire suppressed habitats.



Roadsides through scrub communities managed with regular mowing are typically sparse of vegetation with much of the sandy soil exposed. This right of way is no exception. Patience will be a necessary component for any management plan that depends on migration of wildflowers from adjacent properties in sand scrub communities.



However, there are several species of showy wildflowers present in the adjacent scrub that, over time, could make this right of way a very attractive display of native flora.

It is the intent of the following management plan to provide a baseline for development of a program of adaptive management. Through careful monitoring and periodic analysis, this plan is expected to undergo modifications as conditions change. This plan will balance the need for roadside safety and the ecological needs of the plant communities by defining and scheduling the maintenance activities of regular roadside vegetation management.

#### A. General Criteria Continually Applicable

- Fertilizers shall not be applied to the right of way. No application of materials including soil, seed, mulch, sod, hay or other organic substances is permitted on the right of way without prior written consent of the District Three Vegetation Management Specialist (D3 VMS).
   Any and all sod, mulch, hay and straw used during maintenance activities must be certified free of noxious weeds and their propagules.
- 2. Prior to mowing activities, all equipment must be cleaned and free of soil, seeds, vegetative matter or other debris that could contain or hold noxious weed propagules.
- All mowing must be performed when the ditch is dry enough to prevent rutting. Alternative use of a mower equipped with a hydraulic boom is permitted. At no time shall the height of mowing be less than six inches.

#### **B.** Management Summary

- Semi-annual surveys will be performed by the D3 VMS to locate and record the occurrence
  of invasive plant species, the occurrence of species of concern including federally and/or
  state listed endangered or threatened species, and the occurrence of plant species that
  present a roadside safety hazard. Special management practices that may include spot
  treatment of herbicides or hand clearing will be performed by the Asset Maintenance
  Contractor (AMC) to remove problem species and protect species of concern.
- T-1 routine roadside mowing shall be scheduled by binding language of the current contract
  of the AMC. In the absence of such language, mowing shall be as described in the FDOT
  Guide for Roadside Vegetation Management and shall occur during the months of April,
  May, June, July, August and September on a thirty day mowing cycle as weather conditions
  permit.
- 3. T-2 full right of way mowing <u>may</u> occur during lightning season (May 15 to July 15) and during dormant season (December 15 to February 15) according to need as authorized in writing by the D3 VMS.
- 4. Roadside mowing heights shall be according to the guidelines of the *FDOT Guide for Roadside Vegetation Management* and the *FDOT Maintenance Rating Program Handbook* or as authorized in writing by the D3 VMS.
- 5. Full right of way (T-2) mowing shall occur from the edge of pavement to the limit of the right of way line or wood line, whichever is less.
- 6. Routine roadside mowing (T-1) shall occur in the 15 feet adjacent to the edge of pavement or to the toe of the front slope, whichever is less.
- 7. Any use of prescribed fire for management purposes must be at the direction of the Florida Forest Service.

Notable wildflower species surveyed in 2016 included:	
Baptisia lanceolata	
Baptisia lecontei	
Chrysopsis spp.	
Conradina canescens	
Croptilon divaricatum	
Dalea pinnata	
Eriogonum tomentosum	
Gaillardia pulchella	
Heterotheca subaxillaris	
Liatris spp.	
Licania michauxii	
Lupinus villosus	
Pityopsis graminifolia	